

610 Midterm 2 review: Classical Mechanics

Jinchen He

Calculus of variations

$$S = \int_{x_1}^{x_2} dx f[y(x), y'(x), x], \quad y(x_1) = y_1, \quad y(x_2) = y_2$$

find the $y_0(x)$ to minimize S .

Set $y(x) = y_0 + \epsilon \eta(x)$, $\epsilon \rightarrow 0$, to make $dS = 0$, we can get the E-L equation.

$$\frac{\partial f}{\partial y} - \frac{d}{dx} \left(\frac{\partial f}{\partial y'} \right) = 0$$

E-L equation for unconstrained motion

We have the action $S = \int dt L$ and $L = T - V$.

Generalized force: $\frac{\partial L}{\partial q_i} = F_i$

Generalized momentum: $\frac{\partial L}{\partial \dot{q}_i} = P_i$

E-L equation: $\frac{d}{dt} P_i = F_i$

E-L equation for constrained motion directly.

$L = T - U$ where U contains only non-constraint forces, is stationary at true path with respect to all path that satisfy the constrain.

Oct. 11: three examples

Noether's Theorem

Consider a system of particles with pair-like interactions, the Lagrangian is

$$L = \frac{1}{2} \sum_i m_i \dot{x}_i^2 + \sum_{i>j} V(x_i - x_j)$$

Let $x_i' = x_i + a$, where $a \rightarrow 0$ is a constant, then we can rewrite L with x_i' and $\dot{x}_i' = \dot{x}_i$.

$$L' = \frac{1}{2} \sum_i m_i (\dot{x}_i')^2 + \sum_{i>j} V(x_i' - x_j')$$

So, L' has the same form as L . The Lagrangian is "form invariant" (covariant) under the transformation.

Conservation of Angular Momentum

The angular momentum $\vec{A} = \vec{r} \times \dot{\vec{r}}$ is conserved, the plane of \vec{r} and $\dot{\vec{r}}$ must be the same plane.

For ϕ , we will get the conservation of angular momentum. For r , replace $\dot{\phi}$ with A , we will get

$$\begin{aligned}\mu\ddot{r} &= -\frac{d}{dr}U_{eff}(r) \\ \mu\dot{r} \cdot \ddot{r} &= -\frac{d}{dr}U_{eff}(r) \cdot \dot{r} \\ \mu\frac{d}{dt}\left(\frac{\dot{r}^2}{2}\right) &= -\frac{d}{dt}U_{eff}(r) \\ \mu\frac{\dot{r}^2}{2} + U_{eff}(r) &= E = \text{const} \\ \frac{dr}{dt} &= \sqrt{\frac{2}{\mu}(E - U_{eff})}\end{aligned}$$

Hamiltonian Dynamics

We define the Hamiltonian: $H = \sum_i p_i \dot{q}_i - L$, then we have

$$\frac{\partial H}{\partial p_i} = \dot{q}_i; \quad \frac{\partial H}{\partial q_i} = -\dot{p}_i$$

Liouville's Theorem

Consider a region in phase space and follow its evolution over time. In general, the shape of the region will change, but the volume remains the same.

Liouville's Equation

With $\frac{d\rho}{dt} = 0$, we have

$$\frac{\partial \rho}{\partial t} + \{\rho, H\} = 0$$

Poincare Recurrence Theorem

Poisson Bracket

$$\{f, g\} = \sum_i \left(\frac{\partial f}{\partial q_i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q_i} \right)$$

properties:

- $\{f, g\} = -\{g, f\}$
- $\{af + bg, h\} = a\{f, h\} + b\{g, h\}$
- $\{fg, h\} = f\{g, h\} + \{f, h\}g$

- Jacobi Identity $\{f, \{g, h\}\} + \{h, \{f, g\}\} + \{g, \{h, f\}\} = 0$
- $\frac{df}{dt} = \{f, H\} + \frac{\partial f}{\partial t}$
- $\{q_i, q_j\} = \{p_i, p_j\} = 0$ and $\{q_i, p_j\} = \delta_{ij}$

Canonical Transformation

$$\begin{aligned}\frac{\partial H}{\partial p_i} &= \dot{q}_i; & \frac{\partial H}{\partial q_i} &= -\dot{p}_i \\ q_i &\rightarrow Q_i(q, p, t) \\ p_i &\rightarrow P_i(q, p, t) \\ H(p, q, t) &\rightarrow K(P, Q, t)\end{aligned}$$

If the transformation preserve

$$\frac{\partial K}{\partial P_i} = \dot{Q}_i; \quad \frac{\partial K}{\partial Q_i} = -\dot{P}_i$$

we call it canonical transformation.

Restricted Canonical Transformation

$$\begin{aligned}H(p, q) &= K(P, Q) \\ \{Q_i, Q_j\} &= \{P_i, P_j\} = 0 \text{ and } \{Q_i, P_j\} = \delta_{ij}\end{aligned}$$

Theorem: The Poisson Bracket is invariant under canonical transform.

Action-Angle variables

Canonical transform: $(p, q) \rightarrow (I, \theta)$ [respectively]

So that to make $H = H(I)$, then we have

$$\dot{\theta} = \frac{\partial H}{\partial I} \text{ and } I \text{ are constants}$$

This choice is always possible for any 1-D system, I is called the "action variable", θ is "angle variable".

Claim: The correct choice of I is $I = \frac{1}{2\pi} \oint pdq$.

Then we have


- $H = H(I) = E$
- $\dot{\theta} = \frac{\partial H}{\partial I} = \frac{dE}{dI} = \omega$
- $t = \frac{d}{dE} \int_{q(0)}^{q(t)} pdq$
- $\theta = \omega t = \frac{d}{dI} \int_{q(0)}^{q(t)} pdq$

Adiabatic Invariants

consider only 1 d.o.f.

$$H = \frac{p^2}{2m} + V(q)$$

Assume motion is bounded, so motion is periodic.



Let the potential depend on some parameter λ , so that $V = V(q, \lambda)$. We wish to explore what happens if λ changes slowly ("adiabatically") with time. For example, we may change the length of a pendulum.

Since H is time-dependent, energy is no longer conserved.

$$\frac{dE}{dt} = \left(\frac{\partial H}{\partial t} \right)_{p,q} = \left(\frac{\partial H}{\partial \lambda} \right)_{p,q} \cdot \frac{d\lambda}{dt}$$


There are specific combinations of E and λ which remain constant as λ is slowly changed. These combinations are called "adiabatic invariants".

Claim: The adiabatic invariant for this system is

$$I = \frac{1}{2\pi} \oint p dq$$

where p is now $p = \sqrt{2m(E(t) - V(q, \lambda(t)))}$.

With the adiabatic assumption, the orbit in the phase space below can be closed.



e.g.

$$H = \frac{p^2}{2m} + \frac{1}{2}k(t)q^2, \text{ } k \text{ varying slowly}$$

We have $I = \frac{E}{\omega} = E \sqrt{\frac{m}{k}}$, with the adiabatic invariant claim, we can get

$$\frac{E(t_1)}{\sqrt{k(t_1)}} = \frac{E(t_2)}{\sqrt{k(t_2)}}$$

END of classical mechanics.